Check for updates

RESEARCH ARTICLE

REVISED Fluorescent reporter of *Caenorhabditis elegans* Parkin: Regulators of its abundance and role in autophagy-lysosomal dynamics

[version 2; peer review: 2 approved]

Roman Vozdek[1], Bingying Wang[2], Kathy H. Li[3], Peter P. Pramstaller[1], Andrew A. Hicks[1], Dengke K. Ma[2]

[1]Institute for Biomedicine, Eurac Research, Affiliated institute of the University of Lübeck, Bolzano, 39100, Italy
[2]Cardiovascular Research Institute and Department of Physiology, University of California San Francisco, San Francisco, CA, 94158, USA
[3]Department of Pharmaceutical Chemistry, University of California San Francisco, San Francisco, CA, 94158, USA

v2 **First published:** 18 Feb 2022, **2**:23
https://doi.org/10.12688/openreseurope.14235.1
Latest published: 15 Sep 2023, **2**:23
https://doi.org/10.12688/openreseurope.14235.2

Abstract

Background: Parkin, which when mutated leads to early-onset Parkinson's disease, acts as an E3 ubiquitin ligase. How Parkin is regulated for selective protein and organelle targeting is not well understood. Here, we used protein interactor and genetic screens in *Caenorhabditis elegans* (*C. elegans)* to identify new regulators of Parkin abundance and showed their impact on autophagy-lysosomal dynamics and alpha-Synuclein processing.
Methods: We generated a transgene encoding mCherry-tagged *C. elegans* Parkin – Parkinson's Disease Related 1 (PDR-1). We performed protein interactor screen using Co-immunoprecipitation followed by mass spectrometry analysis to identify putative interacting partners of PDR-1. Ribonucleic acid interference (RNAi) screen and an unbiased mutagenesis screen were used to identify genes regulating PDR-1 abundance. Confocal microscopy was used for the identification of the subcellular localization of PDR-1 and alpha-Synuclein processing.
Results: We show that the *mCherry::pdr-1* transgene rescues the mitochondrial phenotype of *pdr-1* mutants and that the expressed PDR-1 reporter is localized in the cytosol with enriched compartmentalization in the autophagy-lysosomal system. We determined that the transgenic overexpression of the PDR-1 reporter, due to inactivated small interfering RNA (siRNA) generation pathway, disrupts autophagy-lysosomal dynamics. From the RNAi screen of putative PDR-1 interactors we found that the inactivated Adenine Nucleotide Translocator *ant-1.1/hANT*, or hybrid ubiquitin genes *ubq-2/hUBA52 and ubl-1/hRPS27A* encoding a single copy of ubiquitin fused to the ribosomal proteins L40 and S27a, respectively, induced PDR-1 abundance and affected lysosomal dynamics. In addition, we

Open Peer Review

Approval Status ✓ ✓

	1	2
version 2 (revision) 15 Sep 2023	✓ view	
version 1 18 Feb 2022	? view	✓ view

1. **Anat Ben-Zvi**, Ben-Gurion University of the Negev, Beersheba, Israel

2. **Prashant Mishra**, UT Southwestern, Dallas, USA

Any reports and responses or comments on the article can be found at the end of the article.

demonstrate that the abundant PDR-1 plays a role in alpha-Synuclein processing.
Conclusions: These data show that the abundant reporter of *C. elegans* Parkin affects the autophagy-lysosomal system together with alpha-Synuclein processing which can help in understanding the pathology in Parkin-related diseases.

Keywords
Parkinson's disease, Parkin, Synuclein, genetic screen, RNA interference, autophagy, C. elegans



This article is included in the Medical Sciences gateway.



This article is included in the Horizon 2020 gateway.



This article is included in the Cell, Molecular and Structural Biology gateway.



This article is included in the Neuroscience gateway.



This article is included in the Marie-Sklodowska-Curie Actions (MSCA) gateway.



This article is included in the Brain Dysfunction and Degeneration collection.

Corresponding author: Roman Vozdek (roman.vozdek@eurac.edu)

Author roles: Vozdek R: Conceptualization, Data Curation, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Wang B**: Investigation; **Li KH**: Formal Analysis, Methodology; **Pramstaller PP**: Funding Acquisition, Writing – Review & Editing; **Hicks AA**: Funding Acquisition, Resources, Writing – Review & Editing; **Ma DK**: Conceptualization, Funding Acquisition, Methodology, Resources, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 844497. This research was also partly funded by the Department of Educational Assistance, University and Research of the Autonomous Province of Bolzano, Italy, through a core funding initiative to the Institute for Biomedicine.
The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2023 Vozdek R *et al*. This is an open access article distributed under the terms of the Creative Commons Attribution License , which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Vozdek R, Wang B, Li KH *et al.* **Fluorescent reporter of *Caenorhabditis elegans* Parkin: Regulators of its abundance and role in autophagy-lysosomal dynamics [version 2; peer review: 2 approved]** Open Research Europe 2023, **2**:23 https://doi.org/10.12688/openreseurope.14235.2

First published: 18 Feb 2022, **2**:23 https://doi.org/10.12688/openreseurope.14235.1

REVISED **Amendments from Version 1**

In the revised version we added functional assay for expressed *mCherry::PDR-1* using assessment of the heteroplasmy of mutated mtDNA (Figure 1B). Also, we further discussed a possibility of artificial biology of the expressed mCherry::PDR-1 from the transgene *dmaIs48*. Specifically, we have further added a notion of differentially expressed *pdr-1* reporter due to the activity of the *rpl-28* promoter or change of function of PDR-1 due to mCherry tag, and added a notion that the abundant *mCherry::pdr-1* might be activity silenced. Nonetheless, we believe that some cellular function of *mCherry::pdr-1* is likely maintained because of its role in mitochondrial morphology and heteroplasmy in *pdr-1* mutants.

Any further responses from the reviewers can be found at the end of the article

Plain language summary

Parkin promotes activity of the cell cleaning machinery by targeting specific damaged proteins and organelles for their degradation, which allows cells, tissues and whole organisms to cope with stress and aging. Individuals carrying genetic mutations in the gene encoding Parkin develop Parkinson's disease at a young age likely due to disrupted processing of the neuronal protein alpha-Synuclein. This phenomenon is evolutionarily conserved because animal models, such as mice, fish or flies, exhibit progressive alpha-Synuclein pathology upon Parkin inactivation. Here, we generated a worm model with fluorescently labeled Parkin, which, when accumulated in the cells, worsened that pathology associated with Parkinson's disease. This study shows that mutations that cause excessive Parkin accumulation in the roundworm *C. elegans* resemble some characteristics of aging-related diseases and could therefore explain the cellular pathology of some Parkin-related diseases, such as Parkinson's disease or cancer.

Introduction

Various cellular responses to stress conditions that can promote either their survival, or death through apoptosis, have evolved to maintain the health of the whole organism. Parkin is an evolutionarily conserved E3-ubiquitin ligase with a broad role in cell stress by targeting various proteins for degradation through ubiquitin-proteasomal system[1]. Several loss-of-function mutations in the human *Parkin RING-in-between-RING (RBR) E3 Ubiquitin Protein Ligase (PRKN)* gene, which encodes Parkin, have been associated with the autosomal recessive juvenile form of Parkinson's disease (PD), a devastating disease due to loss of dopaminergic neurons in the midbrain Substantia Nigra[2]. Since disease-causing mutations are spread across all exons, and may thus affect several aspects of Parkin biology, including autoinhibition, translocation, interaction or ligase activity and its regulation, it raises the possibility of distinct pathogenetic mechanisms induced by various Parkin mutants that occur in different domains of the protein[3]. Indeed, some mutations, such as R275W, appear to have gain-of-function neurotoxic properties with dominant inheritance highlighting the fact that sometimes heterozygous carriers may develop PD[4].

Parkin is expressed in the cytosol and possesses several regulatory and functional domains. The N-terminal part of the protein encodes an autoinhibitory ubiquitin-like domain (Ubl), that structurally resembles ubiquitin (Ub), which is followed by four zinc-coordinating Really Interesting New Gene (RING)-like domains (RING0, RING1, IBR and RING2) that mediate ubiquitin ligase activity[5]. How Parkin is regulated for selective protein targeting is not well understood. Parkin is autoinhibited under basal conditions and requires activation that has been described for its role in mitophagy (autophagy of the damaged mitochondria)[6]. Phosphatase And Tensin Homolog Induced Putative Kinase 1 (PINK1), a sensor of the depolarized mitochondria, is a direct activator of Parkin[3]. Insufficiently imported PINK1 into mitochondria phosphorylates both the Ubl domain of Parkin as well as Ubiquitin at the equivalent residue Serine 65[7]. Both phosphorylation events trigger a binding switch between the Ubl domain and Ub that releases autoinhibition through opening of the E2-binding site in RING1 and the catalytic site in RING2[8]. Activated Parkin is then recruited onto the outer mitochondrial membrane where it mediates ubiquitination of proteins to trigger mitophagy[9].

Functional *C. elegans* Parkin ortholog, that structurally resembles all Parkin domains, including regulatory sites targeted by PINK1 and phospho-Ubiquitin, is encoded by the *pdr-1* gene[10] and regulates accumulation of mitochondrial DNA mutations, mitochondrial fusions as well as the mitochondrial untranslated protein response[11–13]. Next to PDR-1's conserved role in mitochondrial quality control, it also targets the small guanine triphosphatases (GTPase) CED-10 (Cell Death Abnormality) to regulate cytoskeletal rearrangements preventing apoptotic cell engulfment[14].

The activation of the Parkin towards non-mitochondrial targets, such as synaptic endophilin and synaptojanin[15,16], endo-lysosomal GTPase Rab7 (Rat Sarcoma Virus-associated binding)[17] or alpha-Synuclein[18] are unclear. Several phosphorylation sites, S-nitrosylation and S-sulfhydration have been detected though the effect of these modification needs further studies[19,20]. Here, we used the genetics of *C. elegans* to seek regulators of *C. elegans* Parkin ortholog PDR-1. We generated a new functional mCherry reporter of PDR-1 and identified regulators of its abundance. We show that abundant PDR-1 reporter impacts autophagy-lysosomal dynamics and alters processing of exogenously expressed human alpha-Synuclein in *C. elegans*.

Methods

C. elegans strains

Animals were maintained under standard procedure with nematode growth media (NGM) plates unless otherwise stated. The *C. elegans* construct for *dmaIs48* transgene was generated by Invitrogen Gateway recombination cloning technology. Specifically, we cloned the entire *pdr-1* gene, which was amplified by direct polymerase chain reaction (PCR) using 5'-agGGAAGTGGCTCGAGTATGTCTGATGAAATCTCTATA-3' and 5'-GGCCGATGCGGAGCTCTTAATTAAACCAATGGTCCCATT-3' primers, into *pDEST-mCherry* vector in frame to

mCherry reporter gene with *unc-54 3'-untranslated region (3'UTR)*. The *pDEST-mCherry::pdr-1-unc-54 3'-UTR* vector was subsequently recombined with pENTRY vector carrying *rpl-28* promoter sequence. Transgenic strains were generated by germline transformation using microinjection technique. Transgenic construct was injected at 50 ng/µl into the Bristol strain N2 (20 individual nematodes were used) and stable extrachromosomal lines of mCherry positive animals were established. Extrachromosomal array was subsequently integrated by ultraviolet (UV) irradiation (5 individual nematodes were used) and the strain carrying *dmaIs48* was 2x outcrossed.

The strains used were as follow: wild isolate N2, *xmSi(mai-2p::mai-2::GFP::mai-2 3'-UTR), xmSi;pdr-1(gk448), xmSi;drp-1(tm1108), xmSi;eat-3(ad426), xmSi;dmaIs48(rpl-28p::mCherry::pdr-1);pdr-1(gk448), dmaIs48;xmSi, dmaIs48;xmSi; rde-1(dma341), dmaIs48;adIs2122(lgg-1p::GFP::lgg-1), dmaIs48;adIs2122; rde-1(dma341), dmaIs58(rpl-28p::lmp-1::GFP), dmaIs48;dmaIs58, dmaIs48;dmaIs58;rde-1(dma341), uonEx1(alpha-Synuclein::YFP), dmaIs48;uonEx1(alpha-Synuclein::YFP),* LB138 (*him-8(e1489);uaDf5/+), uaDf5/+;pdr-1(gk448); uaDf5/+;dmaIs48, uaDf5/+;dmaIs48;pdr1(gk448), uaDf5/+;dmaIs48; rde-1(ne300), uaDf5/+;dmaIs48; rde-1(ne300);pdr-1(gk448).*

Imaging and image analysis

Animals were mounted onto a 2% agarose pad containing 10 mM sodium azide and imaged with an EVOS FL auto digital microscope for epifluorescence imaging or a confocal LeicaSP8-X confocal laser scanning microscope within 2–5 minutes. For anoxia stress assay, animals were placed into hypoxia incubator chamber (StemCell, 27310) with constant nitrogen flow delivery to achieve nearly 0% oxygen for 6 or 24 hours prior the imaging[21]. For Carbonyl cyanide p-trifluoro-methoxyphenyl hydrazone (FCCP) assay, animals were exposed to 10uM FCCP 1 hour before imaging. ImageJ (Fiji) was used for the quantification of the fluorescent signal and image processing. Identical setting and conditions were used to compare experimental groups with controls. For the quantification of alpha-Synuclein inclusions, we measured size of the fluorescent foci in the body wall muscle cells from the confocal images using Fiji. At least three images representing each condition from three independent biological replicates were analyzed.

Assessment of mtDNA heteroplasmy

The mtDNA heteroplasmy was assessed by quantitative PCR using previously established protocol with a few modifications[22]. Specifically, 20 L4 animals of each biological group were collected and washed in M9 buffer (Na2HPO4 5.8 g/L, KH2PO4 3 g/L, NaCl 0.5 g/L and NH4Cl 1 g/L) and total DNA was extracted by nematode lysis using Proteinase K at 65C for 60 minutes followed by incubation for 10 minutes at 95C. Quantitative PCR was performed on a CFX96 qPCR/Real-Time PCR Module w/ C1000 Touch Thermal Cycler (Bio-Rad) using All-in-One™ qPCR Mix (GeneCopoeia) and the following set of primers to detect truncated mtDNA (5'TGAGACTTTTAATTATTTACATCCC and 5'CAGTGCATTGACCTAGTCATC), wt mtDNA (5'TGAGACTTTTAATTATTTACATCCC and 5'CAATTTTGCGTGCTATTCC) or total mtDNA (5'CTTTAGGTGGGTTGACAGG and 5'GTAACACCCGTGAAAATCC). The threshold cycle (CT) values were obtained from CFX Manager software (Bio-Rad). The average CT (threshold cycle) of triplicate and or duplicate values obtained for each biological group of mtDNA molecules was normalized using the 2–ΔΔCT method[23], relative to the average CT values obtained for the total mtDNA and/or wt mtDNA molecules from the same sample. At least three independent biological experiments were used to determine the normalized CT values of each strain.

Forward genetic screen

Forward genetic screen for altered *pdr-1* reporter mutants after ethyl methanesulfonate (EMS)-induced random mutagenesis was performed as described previously[24,25]. To screen for mutations that alter expression pattern of the mCherry::PDR-1 reporter, we mutagenized 120 L4 animals carrying the *dmaIs48; xmSi[mai-2::GFP]* with 50 mM EMS in M9 for four hours at 20°C with constant rotation. Worms were subsequently washed in M9 and placed on new NGM plates. We observed the F2 progeny for phenotype using a fluorescence dissecting microscope (Nikon SMZ800N). Animals with constitutively bright mCherry fluorescence were isolated and subsequently sequenced by whole-genome sequencing to obtain lists of candidate genes. Whole genome sequencing was carried out at the UCSF Genomics Facility. DNA sequences from each mutant were analyzed using the CloudMap pipeline Unmapped Mutant Workflow[26] available on the open-source Galaxy web-based platform[27]. Ribonucleic acid interference (RNAi) directed against genes with putative causal mutations was performed to confirm phenocopying.

RNA interference

Feeding RNAi was performed as previously described[28]. Five gravid animals carrying *dmaIs48* transgene were placed on NGM media containing ampicillin 25 µg/ml and 1mM Isopropyl β-d-1-thiogalactopyranoside (IPTG) and seeded with bacteria producing the desired double stranded RNA (dsRNA). Progeny were subsequently grown at 23°C and screened for the phenotype in young adult stage. To induce mild RNAi against *ant-1.1*, *ubq-2* and *ubl-1*, that induced larval arrest in the screened progeny under standard procedure, L1 and L2 animals were placed on RNAi plates and screened for the phenotype in young adult stage. Visual examination of the animals was done on fluorescent stereoscope (Nikon SMZ800N). The bacterial clones were obtained from *C. elegans* RNAi collection - Ahringer (Source Bioscience, 3318).

Western blot analysis

Animals were lysed in the Laemmli sample buffer (Bio-Rad, 1610747) supplemented with the reducing agent β-mercaptoethanol, followed by boiling the samples for 10 min. The worm lysates were separated on 4–15% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS PAGE) (Bio-Rad, 4561086). The proteins were transferred to a nitrocellulose membrane (Bio-Rad, 1620167) and subsequently detected by anti-mCherry rat monoclonal antibody (ThermoFisher Scientific, M11217 1:1000), anti-Synuclein Mouse monoclonal antibody (Abnova, MAB5383

1:2000) and re-incubated with anti-histone H3 rabbit polyclonal antibody (Abcam, ab1791 1:2000) as a loading control.

Co-Immunoprecipitation (Co-IP) Analyses

Whole-animal extracts (*dmaIs48; xmSi(mai-2::GFP); rde-1(dma341)*) were prepared by sonication in M9 buffer (22 mM KH2PO4, 42 mM Na2HPO4, 86 mM NaCl) with protease/phosphatase inhibitor mixture (Sigma-Aldrich, 11836153001 1x). Centrifugation-cleared animal extracts were precleaned by control magnetic beads (bmab-20, ChromoTek) for 30 mins at 4°C, and followed by immunoprecipitation with RFP-trap magnetic beads (ChromoTek, rtma-10) at room temperature for one hour. The RFP-trap beads were subsequently washed five times with M9 buffer. Co-immunoprecipitated proteins were eluted by heating at 70 °C in Laemmli buffer (Bio-Rad, 1610747) for 15 min. Eluates were then subjected to SDS-PAGE and stained with Coomassie Blue to visualize the bands for subsequent in-gel digestion.

Protein identification using liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS)

LC-MS/MS analysis following an in-gel digestion was performed by International Research Resource Center in Biomolecular Mass Spectrometry and Proteomics at University of California San Francisco (UCSF). The in-gel digestion was carried out using their standard sample preparation procedure, available here. The proteins in each gel band were reduced, alkylated, and finally digested overnight with 100 ng of Trypsin (Promega, v511c). The resultant peptide mixture was desalted with µC18-ZipTips (Millipore, ZTC18M960), speed vacuum dried, suspended in 0.1% formic acid, and analyzed on a Velos Pro Elite Orbitrap Mass Spectrometer. The mass spectrometric data was obtained in a data-dependent acquisition mode. The data was then converted into peak lists with PAVA[29], a software developed by International Research Resource Center in Biomolecular Mass Spectrometry and Proteomics. The raw data can be also processed by software package MaxQuant. Using ProteinProspector search engine (v5.20.1), the peak lists were searched against the SwissProt *C. elegans* database and the proteins detected by the LC-MS/MS were subsequently identified.

Results

C. elegans Parkin reporter colocalizes with autophagy-lysosomal compartments

To study regulatory pathways of the Parkin we first investigated the mitochondrial morphology in *pdr-1* mutant animals using mitochondrial reporter *xmSi* encoding Mitochondrial ATPase Inhibitor fused to Green Fluorescent Protein (MAI-2::GFP)[30]. We focused on the hypodermal tissue in adult animals, the largest multinucleated syncytial cell in *C. elegans*. We found that *pdr-1* deficiency leads to elongation of the mitochondrial pattern as seen previously using different mitochondrial reporters[12,13]. We compared such morphology with other known mediators of the mitochondrial dynamics. First, we examined the mitochondrial morphology in the genetic mutants of *drp-1* (Dynamin-Related Protein) encoding the ortholog of DRP1 whose activity is required for the mitofission and *eat-3* (Eating: abnormal pharyngeal pumping) encoding ortholog of Optic atrophy 1 (OPA1) whose activity is required for mitofusion. As expected, while *drp-1* mutants exhibited large disorganized mitochondrial stains, the *eat-3* mutants exhibited fragmented mitochondrial pattern. Next, we investigated mitochondrial morphology in animals subjected to anoxic conditions and mitochondrial oxidative phosphorylation uncoupler FCCP (Carbonyl cyanide p-trifluoro-methoxyphenyl hydrazone). Interestingly, animals exposed to six hours of anoxia (near to 0% oxygen) exhibited donut-like shapes while animals exposed to 24 hours of anoxia exhibited hypermitofusion characterized by the enlarged mitochondria of various shapes. On the other hand, the FCCP triggered fragmented round shape morphology, that differs from the mitochondrial fragmentation in *eat-3* mutants, highlighting mitochondrial shape under disrupted membrane potential. These data show that induced mitochondrial elongations due to loss of *pdr-1*/Parkin differ from the anoxia-induced mitochondrial fusions as well as from disrupted fragmentation in *drp-1* mutants (Figure 1A).

We used the *pdr-1* phenotype to validate the newly generated mCherry reporter of PDR-1. We cloned the entire *pdr-1* gene in frame behind the mCherry reporter under ubiquitous promoter *rpl-28*. Using confocal microscopy we found that the newly generated transgenic strain carrying *dmaIs48(rpl-28p::mCherry::pdr-1)* exhibits bright fluorescent foci of mCherry::PDR-1 that do not colocalize with GFP reporter for mitochondria MAI-2::GFP, though still reduced mitochondrial fusions in *pdr-1* mutants (Figure 1A, 1B). We further investigated the functionality of the mCherry::PDR-1 towards mitochondrial biology by exploring the heteroplasmy of mutated mtDNA using LB138 strain carrying *uaDf5/+*, a heteroplasmic ~ 3 kb mtDNA truncation[22,31,32]. We determined that the mCherry::PDR-1 expression significantly reduced levels of truncated mtDNA in WT as well as *pdr-1* mutants (Figure 1B).

Since mCherry::PDR-1 appears to be a functional Parkin reporter towards mitochondrial biology, we investigated whether the mCherry::PDR-1 reporter recruits onto mitochondria upon exposure to FCCP by following the PDR-1 subcellular localization for 30 minutes after the exposure to FCCP. Although the FCCP altered the mitochondrial pattern that formed round shape morphology, we did not observe recruitment of the mCherry::PDR-1 onto mitochondria despite its dynamic expression pattern in both standard and stressed conditions; see Figure 1A, and Supplementary Video 1 and 2 in *Extended data*[33]. To determine the subcellular localization of the mCherry::PDR-1 foci we crossed the *dmaIs48* transgene with the GFP reporter for autophagosome *adIs2122(lgg-1p::GFP::lgg-1)*[34] and GFP(Venus) reporter for lysosome *dmaIs58(rpl-28p::lmp-1::Venus)*[35]. The confocal microscopy showed that the mCherry foci colocalize with both (LC3, GABARAP and GATE-16 family) and LMP-1 (LAMP (lysosome-associated membrane protein) homolog) reporters indicating that mCherry::PDR-1 is targeted to autophagosome-lysosome compartments (Figure 1C). The full data associated with the results are available in *Underlying data*[33].

Endogenous RNAi pathway regulates expression of mCherry::PDR-1

Because of the striking mCherry::PDR-1 expression pattern in *dmaIs48* animals we aimed to identify novel regulators of



Figure 1. The mCherry reporter of *C. elegans* Parkin (PDR-1) is enriched in the autophagy-lysosomal system. A. Exemplar green fluorescent protein (GFP) fluorescence confocal images showing mitochondrial morphology in the hypoderm of *xmSi(mai-2::GFP)* animals under several environmental and genetic stresses, including *drp-1(tm1108), eat-3(ad426) and pdr-1(gk448).* The PDR-1 reporter expressed from *dmaIs48* transgene restores mitochondrial pattern in *pdr-1(gk448)* mutants. **B.** Normalized levels of mutated mtDNA in indicated strains carrying uaDf5/+. Data are presented as mean ± standard deviation (SD) with p-values calculated by Wilcoxon Mann–Whitney rank sum test to compare two independent populations, * $p<0.05$. **C.** Confocal GFP and mCherry fluorescence images showing expression pattern of the mCherry::PDR-1 together with mitochondrial reporter *xmSi(mai-2::GFP),* autophagosomal reporter *adIs2122(lgg-1::GFP)* and lysosomal reporter *dmaIs58(lmp-1::Venus)* in the hypoderm. The PDR-1 reporter colocalizes with the autophagosomal and lysosomal reporters.

the PDR-1 subcellular expression by unbiased approaches. We performed a forward genetic screen using EMS mutagenesis to isolate mutants with an altered expression pattern. From a screen of approximately 100,000 haploid genomes of the *dmaIs48; xmSi(mai-2::GFP)* strain we have isolated 10 independent mutants with 100% penetrance of markedly increased red fluorescent signal while the GFP signal was not altered (Table 1, Figure 2A). Confocal microscopy showed that the mutants exhibit

Table 1. Ethyl methanesulfonate (EMS)-derived mutants with abundant Parkin reporter mCherry:: PDR-1.

mutation	chr.	penetrance	Gene	Protein change	Annotation
dma334	I	100%	*rde-10*	splicing	RNA interference defective protein
dma330	I	100%	*mut-2*	W164STOP	Mutator
dma337	I	100%	*rrf-1*	Q52STOP	RNA-dependent RNA polymerase
dma338	I	100%	*rrf-1*	A944T	RNA-dependent RNA polymerase
dma339	I	100%	*rrf-1*	G648R	RNA-dependent RNA polymerase
dma330	I	100%	*rrf-1*	R1256STOP	RNA-dependent RNA polymerase
dma327	I	100%	*ego-1*	D1305N	RNA-dependent RNA polymerase
dma333	I	100%	*mut-16*	Q808STOP	Mutator
dma341	V	100%	*rde-1*	P963L	Component of RNA-induced silencing complex
dma325	V	100%	*rde-1*	splicing	Component of RNA-induced silencing complex



Figure 2 . The abundant *C. elegans* Parkin (PDR-1) reporter in ethyl methanesulfonate (EMS)-derived mutants. A. Exemplar green fluorescent protein (GFP) (left) and mCherry (right) fluorescence images showing mCherry::PDR-1 and MAI-2::GFP expression in WT and *rde-1(dma341)* mutant. Only mCherry::PDR-1 expression is increased in *rde-1* mutants. **B.** Fluorescence confocal images of the hypoderm showing the expression pattern of mCherry::PDR-1 in WT and *rde-1(dma341)* mutant. **C.** Confocal view of merged MAI-2::GFP and mCherry:: PDR-1 fluorescence in intestine (upper) and hypoderm (lower) of WT and *rde-1(dma341)* mutant. **D.** The mCherry::PDR-1 protein levels determined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS PAGE) followed by western blot analysis using anti-mCherry Ab and anti-histone H3 Ab. *C. elegans* protein lysates from various genetic backgrounds were analyzed. 50 µg protein samples were loaded per lane. WT represents parental strain carrying *dmaIs48(mCherry::pdr-1)*. Analyzed mutants with *dmaIs48* are as follows: 1. *rde-1(dma325),* 2. *ego-1(dma327)*, 3. *rrf-1(dma339)*, 4. *rde-1(dma341)*, 5. *mut-2(dma340).* N2 represents wild isolate N2.

increased fluorescent signal in the cytosol as well as increased number and size of the foci (Figure 2B) and revealed accumulated mCherry::PDR-1 reporter across the tissues with a strong signal in intestine and hypoderm (Figure 2C). Interestingly, the expression pattern of accumulated mCherry::PDR-1 in the EMS-derived mutants exhibited reduced dynamics in the hypodermal tissue; see Supplementary Video 3 in *Extended data*[33]. Western-Blot analysis using an anti-mCherry antibody showed increased PDR-1 reporter levels in the mutants accompanied with partial processing of the PDR-1 reporter, which is indicated by detection of the shorter variants, compared to parental strain (Figure 2D). Whole genome sequencing of the isolated mutants revealed that 4 mutants carry the protein-changing mutations in the *rrf-1* gene. The *rrf-1* encodes RNA-directed RNA polymerase and is a component of the WAGO 22G RNAs RNAi pathway that generate siRNA for transcriptional silencing of exogenous RNAs[36]. Notably, all the remaining isolated mutants carry at least one protein-changing mutation in the genes of the RNAi pathway, such as *rde-1*, *rde-10*, *ego-1*, *mut-2* and *mut-16* (Table 1). To verify causality of these mutations we employed RNAi against *rrf-1*, *rde-1* and *mut-16* using parental strain and, indeed, we observed that all three RNAi conditions increased reporter levels and thus phenocopied isolated mutants. These data show that increased levels of the mCherry::PDR-1 in the isolated EMS-derived mutants are due to loss of transgene silencing activity at the post-transcriptional level. To evaluate functionality of the abundant mCherry::PDR-1 we used loss-of-function allele *rde-1(ne300)* to induce expression of mCherry::PDR-1 and assessed the heteroplasmy of truncated mtDNA. We determined that the overexpressed mCherry::PDR-1 did not significantly reduce levels of truncated mtDNA neither in WT nor *pdr-1* mutants suggesting that the abundant mCherry::PDR-1 might be activity silenced and may induce artificial cellular responses (Figure 1A). The full data associated with the results are available in *Underlying data*[33].

Hybrid ubiquitins and mitochondrial ANT and ATPase regulate mCherry::PDR-1 levels

Although unbiased mutagenesis revealed regulators of the transgenic mCherry::PDR-1 expression, novel alleles regulating PDR-1 function have not been isolated. Therefore, we employed Co-immunoprecipitation (CoIP) using RFP-trap to identify putative interacting proteins with the mCherry::PDR-1 reporter. We used crude extract of both parental strain *dmaIs48* as well as isolated mutant *dmaIs48; rde-1(dma341)* as the baits. One-dimensional sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS PAGE) revealed five bands of protein pull down that did not differ in size between parental and mutant samples (Figure 3A). We analyzed these bands by gel digest followed by liquid chromatography-tandem mass spectrometry (GeLC-MS/MS) analysis and identified putative interactors with the highest score for adenosine triphosphate (ATP)-Citrate Lyase ACLY-1/hACLY, Heat Shock 70kDa HSP-1/hHSPA1 and hHSPA8,



B **RNAi screen of PDR-1 interactors**

Gene	Ortholog	RNAi
acly-1	ACLY	+
acly-2	ACLY	++
hsp-1	HSPA8	+*
atp-1	ATP5F1A	+++*
atp-2	ATP5F1B	+++*
hsp-60	HSPD1	+
ant-1.1	ANT	+++*
par-5	14-3-3 zeta	+
ftt-2	14-3-3 zeta	+
phb-1	PHB	+
vdac-1	VDAC1	+
vha-8	ATP6V1E1	+*
rps-19	RPS19	+

***dmaIs48* regulators**

Gene	Ortholog	RNAi
rrf-1		+++++
rde-1		+++++
mut-16		+++++

genes with a connection to Parkin function

function	Gene	Ortholog	RNAi
Mitochondrial dynamics	*eat-3*	OPA1	+
	drp-1	DRP1	+
mitophagy	*pink-1*	PINK1	+
	tin-44	TIM-44	+
autophagy	*atg-5*	ATG5	+
	lgg-1	ATG8	+
	lgg-2	ATG8	+
	sqst-1	p62	+
	lrk-1	LRRK2	++
lysosome	*lmp-1*	LAMP	+
PDR-1::GFP interactors	*ubc-2*	E2	+
	ubc-15	E2	+
	ubc-18	E2	+
	chn-1	ChIP	+
	rpt-2	PSMC1	+
ubiquitins	*ubq-1*	UBC	+*
	ubq-2	UBA52	+++++*
	ubl-1	RPS27A	+++++
	ubl-5	UBL5	+
	ned-8	NEDD8	+

Figure 3. Protein interactor and RNAi screen identify regulators of *C. elegans* Parkin (PDR-1) abundance. A. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS PAGE) followed by Coomassie staining of mCherry::PDR-1 pull down. Red fluorescent protein (RFP)-trap was used for Co-immunoprecipitation of protein extracts from *dmaIs48(mCherry::pdr-1)* and *dmaIs48;rde-1(dma341)* animals. Mass spectrometry (MS) analysis of individual bands was used to identify putative interactors. Proteins with the highest peptide coverage are indicated. **B.** Ribonucleic acid interference (RNAi) screen of indicated genes for abundant mCherry::PDR-1.The "plus" symbol indicates mCherry fluorescence intensity and the "star" symbol indicates larval arrest phenotype based on visual examination.

14-3-3-like proteins PAR-5/hYWHAZ and FTT-2/hYWHAZ and several Ribosomal Proteins. Notably, subunits of the Vacuolar ATPase (VHA-8, VHA-12, VHA-13), and several mitochondrial proteins, such as mitochondrial Adenine Nucleotide Translocator ANT-1.1/hANT, ATP synthase subunits ATP-1/hATP5F1A and ATP-2 hATP5F1B, Voltage Dependent Anion Channel VDAC-1/hVDAC1, or Heat Shock Protein HSP-60/hHSPD1, have been also identified; see Supplementary Table 1 in *Extended data*[33].

Next, we performed reverse genetic screen by RNAi against putative PDR-1 regulators. First, we have systematically screened the genes encoding the identified PDR-1 interactors for altered expression pattern of the mCherry::PDR-1 reporter. We found that mCherry::PDR-1 abundance was enhanced by RNAi against mitochondrial *ant-1.1*, *atp-1* and *atp-2* accompanying with developmental arrest at early stages of larva development. Because ANT has been recently identified driver of mitophagy trough Parkin activity[37], we hypothesized that impaired mitophagy may lead to Parkin accumulation. However, the RNAi against *pink-1*/hPINK1 and *tin-44*/hTIM44 (Transport to Inner Mitochondrial Membrane (yeast TIM)), the components of the ANT/Parkin-mediated mitophagy did not increase the expression of mCherry::PDR-1. We have employed RNAi against other genes with a connection to Parkin function, including previously identified PDR-1::GFP interactors[10] or genes encoding ubiquitin, and found that inactivation of hybrid ubiquitin genes *ubl-1* (Ubiquitin-like) and *ubq-2* (ubiquitin) increased the abundance of mCherry::PDR-1, accompanying with developmental arrest at early stages of larva development. These data suggest that accumulation of the PDR-1 reporter is not related to impaired initiation of mitophagy but rather due to impaired mitochondrial function, ubiquitination or proteasomal degradation. The full data associated with the results are available in *Underlying data*[33].

Accumulated mCherry::PDR-1 disrupts lysosomal dynamics

Since increased expression of the PDR-1 reporter in the *rde-1* mutants lost its dynamic pattern in the hypoderm we asked whether its subcellular localization differ compared to the parental strain. We crossed the *mCherry::pdr-1* reporter with autophagy *lgg-1::GFP* and hypodermal *lmp-1::Venus* reporters and found that accumulated PDR-1:mCherry in the *rde-1* mutants does not colocalize with LGG-1:GFP reporter, though the *ant-1.1* and *ubl-1* knockdowns upon mild RNAi still showed partial colocalization (Figure 4A). Interestingly, all conditions, including *rde-1(dma341),* RNAi *ant-1.1* and RNAi *ubl-1*, induced expression of both mCherry::PDR-1 and LMP-1::Venus reporters when expressed together, and colocalized in the lysosomal-like structures. Notably, the dynamics of the PDR-1 reporter in *ubl-1* knockdowns was interrupted (Supplementary Video 4, *Extended data*[33]) compared to control and the LMP-1::Venus expression was increased at the plasma membrane where mCherry::PDR-1 has not been detected (Figure 4A). Because the *lmp-1::Venus* transgene also uses the same ubiquitous promotor *rpl-28* as *mCherry::PDR-1*, we asked whether accumulated LMP-1::Venus in animals with accumulated mCherry::PDR-1 is dependent on the co-expressed *mCherry::pdr-1*. We employed RNAi against *mut-16* and *ubq-2*, the most striking RNAi conditions that induce mCherry::PDR-1 accumulation, in the animals carrying only the *lmp-1::Venus* transgene. We found that expression of LMP-1::Venus was not induced and retained its dynamics in respective knockdowns indicating that significant accumulation of the LMP-1::Venus in *mut-16* and *ubq-2* knockdowns is dependent on the co-expression of the mCherry::PDR-1 (Figure 5A and 5B). These data indicate that lysosomal dynamics are disrupted due to the abundant mCherry::PDR-1.The full data associated with the results are available in *Underlying data*[33].

The mCherry::PDR-1 alters formation of alpha-Synuclein inclusions

We then questioned whether disrupted autophagy-lysosomal dynamics in animals with accumulated mCherry::PDR-1 in cytosol and lysosomes alters proteostasis in the *C. elegans* model of the Parkinson's disease. We used a previously well-established GFP-based reporter of alpha-Synuclein aggregation in body wall muscle cells *uonEx1(unc-54p::alpha-Synuclein::YFP)* and with confocal microscopy quantified the size of the YFP fluorescent foci in the presence and absence of the overexpressed mCherry::PDR-1. First, we determined that the size of the foci (inclusions of aggregated signal) was reduced in the young adult animals co-expressing mCherry::PDR-1. In addition, fluorescent microscopy together with western blot analysis revealed that the expression levels of the alpha-Synuclein::YFP were reduced in the presence of the mCherry::PDR-1 (Figure 6). Next, we asked whether animals with accumulated mCherry::PDR-1, such as *mut-16* and *ubq-2* knockdowns, would further reduce expression of the alpha-Synuclein::YFP. We found that RNAi against *mut-16* reduced expression levels of alpha-Synuclein::YFP in the absence as well as in the presence of the mCherry::PDR-1 accompanied with reduction of the YFP foci. These data suggest that reduction of formed alpha-Synuclein::YFP foci is a result of its reduced expression levels. Interestingly, the RNAi against *ubq-2* increased expression levels of alpha-Synuclein::YFP in the absence of the mCherry::PDR-1, while the *ubq-2* knockdowns with accumulated mCherry::PDR-1 exhibited significantly increased formation of the alpha-Synuclein::YFP foci despite its reduced expression levels (Figure 6). These data indicate that accumulated mCherry::PDR-1 due to inactivated *ubq-2* alters cellular processing of exogenously expressed alpha-Synuclein. The full data associated with the results are available in *Underlying data*[33].

Discussion

Parkin has an active role during proteotoxic as well as mitochondrial stress. Understanding the regulatory mechanisms distinguishing specific Parkin targets may explain its role in Parkinson's disease, tumorigenesis, and other Parkin-associated diseases. Here we present a new functional mCherry reporter of *C. elegans* Parkin (PDR-1) that exhibits autophagy-lysosomal targeting and disrupts the dynamic movements of this system upon its intracellular accumulation. We demonstrate that the *mCherry::pdr-1* transgene reduced mitochondrial fusion in



Figure 4. Abundant *C. elegans* Parkin (PDR-1) reporter alters lysosomal morphology. A. Enlarged view of confocal green fluorescent protein (GFP) and mCherry fluorescence images showing expression pattern of the autophagosomal reporter *adIs2122(lgg-1::GFP)* and PDR-1 reporter *dmaIs48(mCherry::pdr-1)* in the hypoderm of indicated mutants/knockdowns. **B.** Enlarged view of confocal GFP and mCherry fluorescence images showing lysosomal reporter *dmaIs58(lmp-1::Venus)* and PDR-1 reporter *dmaIs48(mCherry::pdr-1)* in the hypoderm of indicated mutants/knockdowns.



Figure 5. Abundant *C. elegans* Parkin (PDR-1) reporter increases expression of lysosomal reporter LMP-1. A. Confocal green fluorescent protein (GFP) and mCherry fluorescence images showing lysosomal reporter *dmaIs58(lmp-1::Venus)* in the absence and presence of the PDR-1 reporter *dmaIs48(mCherry::pdr-1)* upon various ribonucleic acid interference (RNAi) assay. **B.** Quantification of the GFP fluorescence intensity detected in the widefield image of animals carrying *dmaIs58(lmp-1::Venus)* in the absence (-) and presence (+) of the *dmaIs48(mCherry::pdr-1)* upon various RNAi assay. 3 animals from independent biological replicates were analysed. Data are presented as mean ± standard deviation (SD) with p-values calculated by one-way analysis of variance (ANOVA) with Bonferroni correction, * p<0.05.

pdr-1 mutants and reduced levels of mutated mtDNA, indicating that the mCherry::PDR-1 reporter retains some functionality towards mitochondrial biology. On the other hand, we cannot exclude the possibility that mCherry tag at the N-terminal autoinhibitory Ubl domain may alter its regulation, targeting and/or subcellular localization. In addition, differently expressed mCherry::*pdr-1* compared to endogenous *pdr-1* gene due to activity of the *rpl-28* promoter as well as overexpressed *mCherry::pdr-1* may induce artificial responses that need to be evaluated in future studies.

Our preliminary protein interactor screen revealed several putative PDR-1 targets, such as VDAC-1 or HSP-1 which are the orthologs of previously identified human Parkin interactors VDAC1 and HSPA8[38,39]. Interestingly, several identified interactors, such as ANT-1.1/hANT and ATP-1/hATP5F1A, are components of the inner mitochondrial membrane, and therefore we suspect that they are targeted by Parkin during the process of mitochondrial turnover, when the outer mitochondrial membrane is ruptured. Congruently, a similar assay with human Parkin identified several mitochondrial proteins whose direct physical interaction is proposed to be taken part during mitophagy after disruption of the mitochondrial membranes[40]. In fact, Parkin-mediated mitophagy requires both the proteasome system for outer- and autophagy system for inner-mitochondrial membrane rupture[41].

From RNAi screen directed against putative PDR-1 interactors we identified hybrid ubiquitin genes *ubq-2* and *ubl-1* or mitochondrial *ant-1.1*, *atp-1 and atp-2* as regulators of the PDR-1 abundance. How exactly these genes maintain mCherry::PDR-1 levels is not clear. Both *ubq-2/hUBA52* and *ubl-1/hRPS27A* encode a single copy of the ubiquitin domain which can be cleaved from the ribosomal subunits L40 and S27a, respectively[42,43]. It is therefore possible that UBL-1 and UBQ-2 may serve as the ubiquitin substrates for the ATP-dependent PDR-1 auto-ubiquitination. The western blot analysis of overexpressed PDR-1 reporter indicated its partial processing of the mCherry::PDR-1 which supports this hypothesis. Notably, we also found that inactivation of the *ubq-2* increased levels of the exogenously expressed alpha-Synuclein reporter in body wall muscle cells while the co-expression of the mCherry::pdr-1 reduced alpha-Synuclein levels indicating a role of UBQ-2 along with expressed PDR-1 reporter in the regulation of alpha-Synuclein expression levels. Congruently, beneficial effect of Parkin on alpha-Synuclein toxicity has been demonstrated in midbrain cell cultures[44] as well as in a *Drosophila* model of Parkinson's disease[45]. Parkin can mediate K63-linked poly-ubiquitination of alpha-Synuclein to trigger its degradation through increased aggregation[46,47]. Whether UBQ-2/hUBA52 serves as a ubiquitin substrate for the PDR-1/hParkin ubiquitin ligase activity targeting alpha-Synuclein merits further studies.



Figure 6. The *C. elegans* Parkin (PDR-1) alters alpha-Synuclein processing. A. Representative confocal green fluorescent protein (GFP) fluorescence images of the body wall muscle cells in animals carrying uonEx1(alpha-Synuclein::YFP) in the absence and presence of the *dmaIs48(mCherry::pdr-1)* upon various ribonucleic acid interference (RNAi) assay. **B** The alpha-Synuclein::YFP protein levels as determined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS PAGE) followed by western blot using anti-alpha-Synuclein Ab (Abnova MAB5383). Twenty young adult animals collected from three biological replicates upon various RNAi were loaded per lane. **C.** Size quantification of detected fluorescent foci in body wall muscle cells expressing alpha-Synuclein::YFP in the absence (-) and presence (+) of the *dmaIs48(mCherry::pdr-1)* upon various RNAi assay. 3 body wall muscle cells from independent biological replicates were analysed. **D.** Exemplar GFP and mCherry fluorescence images of animals carrying *uonEx1(alpha-Synuclein::YFP)* in the absence and presence of the *dmaIs48(mCherry::pdr-1)* upon indicated RNAi assay. Arrows indicate animals carrying *dmaIs48(mCherry::pdr-1)* which exhibit reduced GFP fluorescence associated with expression levels of alpha-Synuclein reporter. Data are presented as mean ± standard deviation (SD) with p-values calculated by one-way analysis of variance (ANOVA) with Bonferroni correction, * $p<0.05$.

Intriguingly, the abundant PDR-1 reporter due to inactivated *ubq*-2 altered also alpha-Synuclein processing characterized by enlarged fluorescent foci. Since the RNAi of mitochondrial and ubiquitin genes, which induce abundance of the PDR-1 reporter, results in the developmental arrest we have not been able to investigate whether the enlarged aggregated alpha-Synuclein signal may be beneficial and less toxic than diffused alpha-Synuclein, as observed in other model systems[48]. Nonetheless, we hypothesize that abundant PDR-1 may sequester intracellular ubiquitin and thus prevent its binding to alternative E3 ubiquitin ligases. Depleted ubiquitin would then result in insufficient ubiquitination of damaged proteins with consequent impairment of the lysosomal dynamics and proteostasis. Interestingly, the abundant PDR-1 along with the affected lysosomal system due to inactivated *mut-16* did not alter alpha-Synuclein processing. We believe that the reduced formation of the aggregated alpha-Synuclein signal in *mut-16* knockdowns is due to reduced expression levels of alpha-Synuclein reporter in these animals.

Interestingly, lysosomal disfunction with altered lysosomal morphology, that resemble lysosomal morphology in *C. elegans* with abundant PDR-1, has been observed also in the Parkin-deficient cell line[49,50] indicating that both inactivated as well as abundant Parkin may affect lysosomal morphology and function. Taken together, since the enlarged inclusions of alpha-Synuclein in our model system resemble Parkinson's disease pathology, we propose that an affected lysosomal system upon PDR-1 abundance may be one of the possible pathological mechanisms of mutated Parkin.

Ethics and consent

In this project we utilize a roundworm *Caenorhabditis elegans* as a genetic model. These microscopic worms are invertebrate and, therefore, they do not fall within the scope of EU Directive 2010/63/EU on the protection of animals used for scientific purposes. No ethical approval was required for this research.

Data availability

Underlying data

Zenodo: *Caenorhabditis elegans* Parkin: regulators of its abundance and role in autophagy-lysosomal dynamics. https://doi.org/10.5281/zenodo.5940065[33].

This project contains the following underlying data:

- List of Underlying Data.docx (contains description of individual underlying data files in each folder).
- Figure 1.zip (The mCherry reporter of C. elegans Parkin (PDR-1) is enriched in the autophagy-lysosomal system).
- Figure 2.zip (The abundant PDR-1 reporter in ethyl methanesulfonate (EMS)-derived mutants).
- Figure 3.zip (Protein interactor and RNAi screen identify regulators of PDR-1 abundance).
- Figure 4.zip (Abundant PDR-1 reporter alters lysosomal morphology).
- Figure 5.zip (Abundant PDR-1 reporter increases expression of lysosomal reporter LMP-1).
- Figure 6.zip (The PDR-1 alters alpha-Synuclein processing).

Extended data

Zenodo: *Caenorhabditis elegans* Parkin: regulators of its abundance and role in autophagy-lysosomal dynamics. https://doi.org/10.5281/zenodo.5940065[33].

This project contains the following extended data:

- List of Underlying Data.docx (contains description of individual extended data files).
- Extended Data.zip (Supplementary Table 1 and Supplementary Videos 1-5).

Data are available under the terms of the Creative Commons Attribution 4.0 International license (CC-BY 4.0).

Acknowledgments

We thank R. E. Navarro lab and *Caenorhabditis* Genetics Center for *C. elegans* strains. Mass Spectrometry was provided by the Mass Spectrometry Resource at UCSF (A.L. Burlingame, Director) supported by the Dr. Miriam and Sheldon G. Adelson Medical Research Foundation (AMRF) and the NIH-NIGMS. R.V. was supported by International Mobility Program of the Autonomous Province of Bolzano.

References

1. Shimura H, Hattori N, Kubo S, *et al.*: **Familial Parkinson disease gene product, parkin, is a ubiquitin-protein ligase.** *Nat Genet.* 2000; **25**(3): 302–5.
PubMed Abstract | Publisher Full Text

2. Nuytemans K, Theuns J, Cruts M, *et al.*: **Genetic etiology of Parkinson disease associated with mutations in the *SNCA, PARK2, PINK1, PARK7*, and *LRRK2* genes: a mutation update.** *Hum Mutat.* 2010; **31**(7): 763–80.
PubMed Abstract | Publisher Full Text | Free Full Text

3. Narendra D, Walker JE, Youle R: **Mitochondrial Quality Control Mediated by PINK1 and Parkin: Links to Parkinsonism.** *Cold Spring Harb Perspect Biol.* 2012; **4**(11): a011338.
PubMed Abstract | Publisher Full Text | Free Full Text

4. Cookson MR, Lockhart PJ, McLendon C, *et al.*: **RING finger 1 mutations in Parkin produce altered localization of the protein.** *Hum Mol Genet.* 2003; **12**(22): 2957–65.
PubMed Abstract | Publisher Full Text

5. Seirafi M, Kozlov G, Gehring K: **Parkin structure and function.** *FEBS J.* 2015;

282(11): 2076–88.
PubMed Abstract | Publisher Full Text | Free Full Text

6. Narendra D, Tanaka A, Suen DF, *et al.*: **Parkin is recruited selectively to impaired mitochondria and promotes their autophagy.** *J Cell Biol.* 2008; **183**(5): 795–803.
PubMed Abstract | Publisher Full Text | Free Full Text

7. Kazlauskaite A, Martínez-Torres RJ, Wilkie S, *et al.*: **Binding to serine 65-phosphorylated ubiquitin primes Parkin for optimal PINK1-dependent phosphorylation and activation.** *EMBO Rep.* 2015; **16**(8): 939–54.
PubMed Abstract | Publisher Full Text | Free Full Text

8. Gladkova C, Maslen SL, Skehel JM, *et al.*: **Mechanism of parkin activation by PINK1.** *Nature.* 2018; **559**(7714): 410–4.
PubMed Abstract | Publisher Full Text | Free Full Text

9. Koyano F, Okatsu K, Kosako H, *et al.*: **Ubiquitin is phosphorylated by PINK1 to activate parkin.** *Nature.* 2014; **510**(7503): 162–6.
PubMed Abstract | Publisher Full Text

10. Springer W, Hoppe T, Schmidt E, *et al.*: **A *Caenorhabditis elegans* Parkin mutant with altered solubility couples α-synuclein aggregation to proteotoxic stress.** *Hum Mol Genet.* 2005; **14**(22): 3407–23.
PubMed Abstract | Publisher Full Text

11. Ahier A, Dai CY, Kirmes I, *et al.*: **PINK1 and parkin shape the organism-wide distribution of a deleterious mitochondrial genome.** *Cell Rep.* 2021; **35**(9): 109203.
PubMed Abstract | Publisher Full Text

12. Luz AL, Rooney JP, Kubik LL, *et al.*: **Mitochondrial Morphology and Fundamental Parameters of the Mitochondrial Respiratory Chain Are Altered in *Caenorhabditis elegans* Strains Deficient in Mitochondrial Dynamics and Homeostasis Processes.** *PLoS One.* 2015; **10**(6): e0130940.
PubMed Abstract | Publisher Full Text | Free Full Text

13. Cooper JF, Machiela E, Dues DJ, *et al.*: **Activation of the mitochondrial unfolded protein response promotes longevity and dopamine neuron survival in Parkinson's disease models.** *Sci Rep.* 2017; **7**(1): 16441.
PubMed Abstract | Publisher Full Text | Free Full Text

14. Cabello J, Sämann J, Gómez-Orte E, *et al.*: **PDR-1/hParkin negatively regulates the phagocytosis of apoptotic cell corpses in *Caenorhabditis elegans*.** *Cell Death Dis.* 2014; **5**(3): e1120.
PubMed Abstract | Publisher Full Text | Free Full Text

15. Trempe JF, Chen CXQ, Grenier K, *et al.*: **SH3 Domains from a Subset of BAR Proteins Define a Ubl-Binding Domain and Implicate Parkin in Synaptic Ubiquitination.** *Mol Cell.* 2009; **36**(6): 1034–47.
PubMed Abstract | Publisher Full Text

16. Cao M, Milosevic I, Giovedi S, *et al.*: **Upregulation of Parkin in Endophilin Mutant Mice.** *J Neurosci.* 2014; **34**(49): 16544–9.
PubMed Abstract | Publisher Full Text | Free Full Text

17. Song P, Trajkovic K, Tsunemi T, *et al.*: **Parkin Modulates Endosomal Organization and Function of the Endo-Lysosomal Pathway.** *J Neurosci.* 2016; **36**(8): 2425–37.
PubMed Abstract | Publisher Full Text | Free Full Text

18. Shimura H, Schlossmacher MG, Hattori N, *et al.*: **Ubiquitination of a new form of α-synuclein by parkin from human brain: implications for Parkinson's disease.** *Science.* 2001; **293**(5528): 263–9.
PubMed Abstract | Publisher Full Text

19. Chung KKK, Thomas B, Li X, *et al.*: **S-Nitrosylation of Parkin Regulates Ubiquitination and Compromises Parkin's Protective Function.** *Science.* 2004; **304**(5675): 1328–31.
PubMed Abstract | Publisher Full Text

20. Vandiver MS, Paul BD, Xu R, *et al.*: **Sulfhydration mediates neuroprotective actions of parkin.** *Nat Commun.* 2013; **4**(1): 1626.
PubMed Abstract | Publisher Full Text | Free Full Text

21. Vozdek R, Long Y, Ma DK: **The receptor tyrosine kinase HIR-1 coordinates HIF-independent responses to hypoxia and extracellular matrix injury.** *Sci Signal.* 2018; **11**(550): eaat0138.
PubMed Abstract | Publisher Full Text

22. Valenci I, Yonai L, Bar-Yaacov D, *et al.*: ***Parkin* modulates heteroplasmy of truncated mtDNA in *Caenorhabditis elegans*.** *Mitochondrion.* 2015; **20**: 64–70.
PubMed Abstract | Publisher Full Text

23. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method.** *Method.* 2001; **25**(4): 402–8.
PubMed Abstract | Publisher Full Text

24. Brenner S: **The genetics of Caenorhabditis elegans.** *Genetics.* 1974; **77**(1): 71–94.
PubMed Abstract | Publisher Full Text | Free Full Text

25. Ma DK, Vozdek R, Bhatla N, *et al.*: **CYSL-1 Interacts with the O_2-Sensing Hydroxylase EGL-9 to Promote H_2S-Modulated Hypoxia-Induced Behavioral Plasticity in *C. elegans*.** *Neuron.* 2012; **73**(5): 925–40.
PubMed Abstract | Publisher Full Text | Free Full Text

26. Minevich G, Park DS, Blankenberg D, *et al.*: **CloudMap: A Cloud-Based Pipeline for Analysis of Mutant Genome Sequences.** *Genetics.* 2012; **192**(4): 1249–69.
PubMed Abstract | Publisher Full Text | Free Full Text

27. Blankenberg D, Kuster GV, Coraor N, *et al.*: **Galaxy: A Web-Based Genome Analysis Tool for Experimentalists.** *Curr Protoc Mol Biol.* 2010; **89**(1): Unit 19.10.1-21.
PubMed Abstract | Publisher Full Text | Free Full Text

28. Kamath RS, Ahringer J: **Genome-wide RNAi screening in *Caenorhabditis elegans*.** *Methods.* 2003; **30**(4): 313–21.
PubMed Abstract | Publisher Full Text

29. Guan S, Price JC, Prusiner SB, *et al.*: **A data processing pipeline for mammalian proteome dynamics studies using stable isotope metabolic labeling.** *Mol Cell Proteomics.* 2011; **10**(12): M111.010728.
PubMed Abstract | Publisher Full Text | Free Full Text

30. Fernández-Cárdenas LP, Villanueva-Chimal E, Salinas LS, *et al.*: ***Caenorhabditis elegans* ATPase inhibitor factor 1 (IF_1) MAI-2 preserves the mitochondrial membrane potential ($\Delta\psi_m$) and is important to induce germ cell apoptosis.** *PLoS One.* 2017; **12**(8): e0181984.
PubMed Abstract | Publisher Full Text | Free Full Text

31. Lin YF, Schulz AM, Pellegrino MW, *et al.*: **Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded protein response.** *Nature.* 2016; **533**(7603): 416–419.
PubMed Abstract | Publisher Full Text | Free Full Text

32. Gitschlag BL, Kirby CS, Samuels DC, *et al.*: **Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in *C. elegans*.** *Cell Metab.* 2016; **24**(1): 91–103.
PubMed Abstract | Publisher Full Text | Free Full Text

33. Vozdek R, Wang B, Li KH, *et al.*: **Caenorhabditis elegans Parkin: regulators of its abundance and role in autophagy-lysosomal dynamics.** *Zenodo.* 2021.
http://www.doi.org/10.5281/zenodo.5940065

34. Kang C, You Y, Avery L: **Dual roles of autophagy in the survival of *Caenorhabditis elegans* during starvation.** *Genes Dev.* 2007; **21**(17): 2161–71.
PubMed Abstract | Publisher Full Text | Free Full Text

35. Luo S, Wang X, Bai M, *et al.*: **The conserved autoimmune-disease risk gene *TMEM39A* regulates lysosome dynamics.** *Proc Natl Acad Sci U S A.* 2021; **118**(6): e2011379118.
PubMed Abstract | Publisher Full Text | Free Full Text

36. Billi AC, Fischer SEJ, Kim JK: **Endogenous RNAi pathways in *C. elegans*.** *WormBook.* 2014; 1–49.
PubMed Abstract | Publisher Full Text | Free Full Text

37. Hoshino A, Wang W, Wada S, *et al.*: **The ADP/ATP translocase drives mitophagy independent of nucleotide exchange.** *Nature.* 2019; **575**(7782): 375–9.
PubMed Abstract | Publisher Full Text | Free Full Text

38. Geisler S, Holmström KM, Skujat D, *et al.*: **PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1.** *Nat Cell Biol.* 2010; **12**(2): 119–31.
PubMed Abstract | Publisher Full Text

39. Imai Y, Soda M, Hatakeyama S, *et al.*: **CHIP Is Associated with Parkin, a Gene Responsible for Familial Parkinson's Disease, and Enhances Its Ubiquitin Ligase Activity.** *Mol Cell.* 2002; **10**(1): 55–67.
PubMed Abstract | Publisher Full Text

40. Zanon A, Rakovic A, Blankenburg H, *et al.*: **Profiling of Parkin-binding partners using tandem affinity purification.** *PLoS One.* 2013; **8**(11): e78648.
PubMed Abstract | Publisher Full Text | Free Full Text

41. Wei Y, Chiang WC, Sumpter R Jr, *et al.*: **Prohibitin 2 Is an Inner Mitochondrial Membrane Mitophagy Receptor.** *Cell.* 2017; **168**(1–2): 224–238.e10.
PubMed Abstract | Publisher Full Text | Free Full Text

42. Komander D, Clague MJ, Urbé S: **Breaking the chains: structure and function of the deubiquitinases.** *Nat Rev Mol Cell Biol.* 2009; **10**(8): 550–63.
PubMed Abstract | Publisher Full Text

43. Kobayashi M, Oshima S, Maeyashiki C, *et al.*: **The ubiquitin hybrid gene *UBA52* regulates ubiquitination of ribosome and sustains embryonic development.** *Sci Rep.* 2016; **6**(1): 36780.
PubMed Abstract | Publisher Full Text | Free Full Text

44. Petrucelli L, O'Farrell C, Lockhart PJ, *et al.*: **Parkin protects against the toxicity associated with mutant α-Synuclein: proteasome dysfunction selectively affects catecholaminergic neurons.** *Neuron.* 2002; **36**(6): 1007–19.
PubMed Abstract | Publisher Full Text

45. Rana A, Rera M, Walker DW: **Parkin overexpression during aging reduces proteotoxicity, alters mitochondrial dynamics, and extends lifespan.** *Proc Natl Acad Sci U S A.* 2013; **110**(21): 8638–43.
PubMed Abstract | Publisher Full Text | Free Full Text

46. Olzmann JA, Li L, Chudaev MV, *et al.*: **Parkin-mediated K63-linked polyubiquitination targets misfolded DJ-1 to aggresomes via binding to HDAC6.** *J Cell Biol.* 2007; **178**(6): 1025–38.
PubMed Abstract | Publisher Full Text | Free Full Text

47. Tan JMM, Wong ESP, Kirkpatrick DS, *et al.*: **Lysine 63-linked ubiquitination promotes the formation and autophagic clearance of protein inclusions associated with neurodegenerative diseases.** *Hum Mol Genet.* 2008; **17**(3): 431–9.
PubMed Abstract | Publisher Full Text

48. Winner B, Jappelli R, Maji SK, *et al.*: ***In vivo* demonstration that α-Synuclein oligomers are toxic.** *Proc Natl Acad Sci U S A.* 2011; **108**(10): 4194–9.
PubMed Abstract | Publisher Full Text | Free Full Text

49. Guerra F, Girolimetti G, Beli R, *et al.*: **Synergistic Effect of Mitochondrial and Lysosomal Dysfunction in Parkinson's Disease.** *Cells.* 2019; **8**(5): 452.
PubMed Abstract | Publisher Full Text | Free Full Text

50. Okarmus J, Bogetofte H, Schmidt SI, *et al.*: **Lysosomal perturbations in human dopaminergic neurons derived from induced pluripotent stem cells with *PARK2* mutation.** *Sci Rep.* 2020; **10**(1): 10278.
PubMed Abstract | Publisher Full Text | Free Full Text

Open Peer Review

Current Peer Review Status: ✔ ✔

Version 2

Reviewer Report 05 October 2023

https://doi.org/10.21956/openreseurope.17936.r35157

© **2023 Ben-Zvi A.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

✔ **Anat Ben-Zvi**
Ben-Gurion University of the Negev, Beersheba, Israel

The authors addressed most of my concerns. The new data suggest that the *mCherry::pdr-1* is active but further indicate that the transgene overexpression modulates its function (*uaDf5* levels in *mCherry::pdr-1* background are 2-fold lower than WT). Thus, the work is clear and technically sound but focuses on the impact of Parkin's overexpression on worms, limiting its applicability to WT PDR-1.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Proteostasis, mitophagy C. elegans

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 29 April 2022

https://doi.org/10.21956/openreseurope.15353.r28892

© **2022 Mishra P.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

✔ **Prashant Mishra**
[1] UT Southwestern, Dallas, USA
[2]

UT Southwestern, Dallas, USA

Vozdek *et al.* report on the use of a mCherry:PDR1 reporter in C.elegans. PDR-1 is an ortholog of mammalian Parkin, which has been implicated in familial forms of Parkinson's disease, mitophagy, and organelle morphology. Here the authors make use of their reporter to identify Parkin interactors in C.elegans, as well as potential functions of Parkin in this model system.

The authors find that their mCheery:PDR1 reporter largely localizes with lysosomal structures, and is not recruited to mitochondria under decoupling conditions (based on image analysis). Mass-spec immunoprecipitation and genetic screens identify a number of mitochondrial proteins as potential interactors and regulators of their PDR1 reporter. In addition, they find that their reporter alters processing of Synuclein inclusions and lysosomal morphology.

Overall the data is convincing and technically sound. The relevance is limited, as the authors are studying an overexpressed and mCherry fused version of Parkin; thus, it is unclear if their findings are relevant to native/endogenous Parkin, particularly since their construct does not replicate key features of endogenous Parkin (including cytosolic localization and mitochondrial recruitment (Fig. 1)). Thus, although the work emphasizes the consequences and potential actions of Parkin overexpression in worms, my major recommendation would be to investigate if these results (interactors, effects on lysosomal dynamics) are properties of endogenous Parkin.

Is the work clearly and accurately presented and does it cite the current literature?
Partly

Is the study design appropriate and does the work have academic merit?
Partly

Are sufficient details of methods and analysis provided to allow replication by others?
Yes

If applicable, is the statistical analysis and its interpretation appropriate?
I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?
Yes

Are the conclusions drawn adequately supported by the results?
Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Mitochondrial Disease; mitochondrial DNA

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 26 April 2022

https://doi.org/10.21956/openreseurope.15353.r28891

© **2022 Ben-Zvi A.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

?

Anat Ben-Zvi
[1] Ben-Gurion University of the Negev, Beersheba, Israel
[2] Ben-Gurion University of the Negev, Beersheba, Israel

The manuscript "Caenorhabditis elegans Parkin: Regulators of its abundance and role in autophagy-lysosomal dynamics" by Vozdek *et al.* aims to identify new regulators of Parkin (pdr-1) in C. elegans. To this end, they establish a novel transgene encoding mCherry-tagged pdr-1. This transgene is enriched in the autophagy-lysosomal system and modulates mitochondrial morphology in the hypoderm. Using this transgene, the authors then perform forward and reverse genetic screens and protein-protein interactions to examine for modifiers of transgene abundance and localization. Finally, they examine the impact of this transgene on alpha-Synuclein inclusions.

The main concern I have is the possibility that the mCherry::PDR-1 construct is toxic, affecting the interpretation of the results. The authors themselves state in the discussion that "we cannot exclude the possibility that mCherry tag at the N-terminal autoinhibitory Ubl domain may alter its regulation, targeting and/or subcellular localization." The authors did not address this concern experimentally. While they showed that mCherry::PDR-1 altered mitochondria morphology (Fig 1A) compared to pdr-1 (mutant?) alone, I do not agree that the construct restored mitochondria morphology to that of WT. Moreover, known (and simple) functional assays examine pdr-1 function and could be used to determine the construct functionality—for example, the impact of pdr-1 on truncated mtDNA levels (see suggested ref below).

The modifiers identified in the forward genetic screen reinforce this concern. Most are associated with RNA interference and silencing and point to the possibility that the expression of the mCherry::PDR-1 construct is activity silenced. In this regard, an important control will be to examine the impact of these mutations on pdr-1(+) expression by examining mRNA, or better yet, protein levels—alternatively, the effect of these mutants on other pdr-1 constructs.

These issues need to be addressed in revisions.

Minor comments:

1. Several works on PDR-1 function in C. elegans are not referenced see the reference list for examples.

2. There is no statistical information in the manuscript. Specifically, Figs. 3B, 5B, and 6C.

3. It is not clear that pdr-1/parkin in the legend refers to pdr-1 mutant and which mutant.

4. What are "lost the dynamic movements in the hypodermal tissue," and how were they measured.

References
1. Lin Y, Schulz A, Pellegrino M, Lu Y, et al.: Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded protein response. *Nature*. 2016; **533** (7603): 416-419 Publisher Full Text
2. Gitschlag B, Kirby C, Samuels D, Gangula R, et al.: Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in C. elegans. *Cell Metabolism*. 2016; **24** (1): 91-103 Publisher Full Text
3. Valenci I, Yonai L, Bar-Yaacov D, Mishmar D, et al.: Parkin modulates heteroplasmy of truncated mtDNA in Caenorhabditis elegans.*Mitochondrion*. 2015; **20**: 64-70 PubMed Abstract | Publisher Full Text

Is the work clearly and accurately presented and does it cite the current literature?
Partly

Is the study design appropriate and does the work have academic merit?
Partly

Are sufficient details of methods and analysis provided to allow replication by others?
Yes

If applicable, is the statistical analysis and its interpretation appropriate?
No

Are all the source data underlying the results available to ensure full reproducibility?
Yes

Are the conclusions drawn adequately supported by the results?
Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: proteostasis, mitophagy C. elegans

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 06 Sep 2023

Roman Vozdek

The manuscript "Caenorhabditis elegans Parkin: Regulators of its abundance and role in autophagy-lysosomal dynamics" by Vozdek et al. aims to identify new regulators of Parkin (pdr-1) in C. elegans. To this end, they establish a novel transgene encoding mCherry-tagged pdr-1. This transgene is enriched in the autophagy-lysosomal system and modulates mitochondrial morphology in the hypoderm. Using this transgene, the authors

then perform forward and reverse genetic screens and protein-protein interactions to examine for modifiers of transgene abundance and localization. Finally, they examine the impact of this transgene on alpha-Synuclein inclusions.

Thank you for your critical reading and constructive comments on our manuscript.
The main concern I have is the possibility that the mCherry::PDR-1 construct is toxic, affecting the interpretation of the results. The authors themselves state in the discussion that "we cannot exclude the possibility that mCherry tag at the N-terminal autoinhibitory Ubl domain may alter its regulation, targeting and/or subcellular localization." The authors did not address this concern experimentally. **We agree that the mCherry::PDR-1 can be toxic for the animals and thus we conclude that the obtained results may be one of the possible pathological mechanisms of mutated Parkin. Because in this study the role of the mCherry tag is not experimentally addressed, we further discuss a possibility of its artificial biology. Specifically, we have further added a notion of differentially expressed pdr-1 due to activity of the *rpl-28* promoter or change of function of PDR-1 due to mCherry (Page 14). Nonetheless, we believe that some cellular function of *pdr-1* is maintained when expressed from *dmaIs48* transgene because of its role in partial restoration of the mitochondrial morphology as well as altered heteroplasmy in *pdr-1* mutants (see below).**

While they showed that mCherry::PDR-1 altered mitochondria morphology (Fig 1A) compared to pdr-1 (mutant?) alone, I do not agree that the construct restored mitochondria morphology to that of WT. Moreover, known (and simple) functional assays examine pdr-1 function and could be used to determine the construct functionality—for example, the impact of pdr-1 on truncated mtDNA levels (see suggested ref below). **We agree with the reviewer that the functional assay of *pdr-1* is needed to further explore the effect of mCherry::PDR-1 on mitochondrial biology. As suggested, we have assessed truncated mtDNA levels in animals expressing *mCherry::pdr-1*. Although we did not determine higher levels of truncated mtDNA in *pdr-1* mutants at statistical significance, we determined that the levels of mutated mtDNA are significantly decreased in WT and *pdr-1* mutants carrying *dmaIs48* transgene. We have added these data into the manuscript (Page 8, Figure 1). In addition, regarding the assessed effect of the transgene on mitochondrial morphology, we have changed the wording to " mCherry::PDR-1 reduced mitochondrial fusions in *pdr-1* mutants." (Page 8). Because of the observed changes of mitochondrial morphology as well as truncated mtDNA levels in *pdr-1* mutants, we revised our conclusion into "mCherry::PDR-1 might retain some functionality towards mitochondria."(Page 8).**

The modifiers identified in the forward genetic screen reinforce this concern. Most are associated with RNA interference and silencing and point to the possibility that the expression of the mCherry::PDR-1 construct is activity silenced. In this regard, an important control will be to examine the impact of these mutations on pdr-1(+) expression by examining mRNA, or better yet, protein levels—alternatively, the effect of these mutants on other pdr-1 constructs.
We have assessed the effect of mutated *rde-1* gene, that causes *mCherry::pdr-1* overexpression, on heteroplasmy of truncated mtDNA and found that the overexpressed *mCherry::pdr-1* lost the capability to significantly reduce truncated

mtDNA. We have revised the conclusion into "These data suggest that the abundant *mCherry::pdr-1* might be activity silenced (Page 9). We agree with the reviewer that the impact of the mutations on endogenous levels of *pdr-1* and different *pdr-1* constructs would be indeed informative, however, we believe that for the conclusions focusing on the role of mCherry::PDR-1 transgene, and not the role of endogenous *pdr-1*, the proposed assays assessing endogenous *pdr-1* and different *pdr-1* reporters would be out of scope of this manuscript. We have changed the title of the manuscript to address this issue.

Minor comments: Several works on PDR-1 function in C. elegans are not referenced see the reference list for examples. **We have added the references. 1. Lin Y, Schulz A, Pellegrino M, et al.: Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded protein response. Nature. 2016; 533 (7603): 416-419 | Publisher Full Text 2. Gitschlag B, Kirby C, Samuels D, et al.: Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in C. elegans. Cell Metabolism. 2016; 24 (1): 91-103 | Publisher Full Text 3. Valenci I, Yonai L, Bar-Yaacov D, et al.: Parkin modulates heteroplasmy of truncated mtDNA in Caenorhabditis elegans. Mitochondrion. 2015; 20 64-70 PubMed Abstract | Publisher Full Text**

There is no statistical information in the manuscript. Specifically, Figs. 3B, 5B, and 6C. **We have added statistical information into 5B and 6C while we do not have exact measures for 3B. We have revised and clarified figure legends.**

It is not clear that pdr-1/parkin in the legend refers to pdr-1 mutant and which mutant. **We have corrected the legend.**

What are "lost the dynamic movements in the hypodermal tissue," and how were they measured. **We have revised this sentence.**

Competing Interests: No competing interests were disclosed.